

Remarks on Bounds for Quantum codes

Alexei Ashikhmin

Los Alamos National Laboratory

Abstract

We present some results that show that bounds from classical coding theory still work in many cases of quantum coding theory.

1 Nondegenerate Quantum Codes

Let Q be an $((n, K, d))$ nondegenerate quantum code with projection operator P . Let A_i and B_i be the quantum enumerators of Q , [1],[2]

$$A_i = \sum_{wt(E)=i} \text{Tr}(EP)\text{Tr}(EP),$$

$$B_i = \sum_{wt(E)=i} \text{Tr}(EPEP),$$

where E is an error operator. (See [1],[2] for definition of error operators and their weights.)

It is shown in [1],[2] that

$$A_t = \frac{1}{2^n} \sum_{i=0}^n B_i P_t(i, n),$$

where $P_t(i, n) = \sum_{j=0}^i (-1)^j 3^{t-j} \binom{i}{j} \binom{n-j}{t-j}$ is the Krawtchouk polynomial. Replace A_i by A_i/K^2 and B_i by B_i/K . Then we get

$$A_t = \frac{1}{2^n K} \sum_{i=0}^n B_i P_t(i, n) \text{ and } A_0 = B_0 = 1. \quad (1)$$

Now we can use well known techniques to get upper bounds for the quantity $2^n K$ (see for example [3]). We describe this technique here for completeness.

Let $f(x) = \sum_{i=0}^n a_i x^i$ be a polynomial . Consider its expansion by Krawtchouk polynomials

$$f(x) = \sum_{i=0}^n f_i P_i(x, n). \quad (2)$$

Then from (1) and (2) we have

$$\sum_{i=0}^n B_i f(i) = \frac{1}{2^n K} \sum_{i=0}^n A_i f_i.$$

Suppose that

$$f_0 > 0, f_i \geq 0, \text{ if } i = 1, \dots, n, \text{ and } f(0) > 0, f(i) \leq 0, \text{ if } i = d, \dots, n. \quad (3)$$

Then taking into account that for nondegenerate codes $A_i = B_i = 0$ for $i = 1, \dots, d-1$, we have

$$2^n K = \frac{f(0) + \sum_{i=1}^n f(i) B_i}{f_0 + \sum_{i=1}^n f_i A_i} \leq \frac{f(0)}{f_0}. \quad (4)$$

From classical coding theory we know many useful polynomials that satisfy conditions (3). Here we list some of them.

Let $f(x) = 4^{q-d+1} \prod_{j=d}^n (1 - \frac{x}{j})$ [4]. Then

$$f_i = \frac{\binom{n-k}{d-1}}{\binom{n}{d-1}} \geq 0.$$

So we get a quantum Singleton bound [5][6]: $K \leq 2^{n-2d+2}$. Note that $f_i = 0$ for $i \geq n-d+2$ and hence if a quantum code meets the Singleton bound it should be nondegenerate up to $n-d+2$ (a quantum code is called nondegenerate up to t if $A_i = 0$ for $i = 1, \dots, t-1$ [6].)(In [4] this proposition is proven for all quantum codes.)

Let $d = 2e + 1$ and

$$f_i = \left(\frac{P_e(i-1, n-1)}{\sum_{i=0}^e 3^i \binom{n}{i}} \right)^2.$$

Then [4] $f(i) = 0$ for $i = d, \dots, n$ and $f(0) = 4^n / \sum_{i=0}^e 3^i \binom{n}{i}$, and from (4) we get the quantum analog of the Hamming bound: $K \leq 2^n \sum_{i=0}^e 3^i \binom{n}{i}$.

Analogously, from [3] it is straightforwardly follows that

$$K \leq L^n(d)/2^n,$$

where

$$L^n(x) = \begin{cases} L_k^n(x), & \text{if } d_k(n-1) + 1 < x < d_{k-1}(n-2) + 1 \\ 4L_k^{n-1}(x), & \text{if } d_k(n-2) + 1 < x < d_k(n-1) + 1. \end{cases}$$

$$L_k^n(x) = \sum_{i=1}^{k-1} \binom{n}{i} 3^i - \binom{n}{k} 3^k \frac{P_{k-1}(x-1, n-1)}{P_k(x, n)},$$

and $d_k(n)$ is the smallest root of $P_k(x, n)$.

Using the asymptotic version of this bound [3], as n tends to infinity

$$\log_2 K \geq 2H_4(x) - n/2 \text{ and } d < n\gamma_4(x), \quad (5)$$

where

$$\gamma_q(x) = 1/q((q-1) - (q-2)x - 2\sqrt{(q-1)x(1-x)}), \text{ and}$$

$$H_q(x) = -x \log_q x - (1-x) \log_q(1-x) + x \log_q(q-1).$$

(See figure 1,B.) Note that as $\frac{\log_2 K}{n}$ tends to zero $\frac{d}{n} \approx 0.316$ which is better then the value $n/3$ from [9]. However the latter bound is valid for all quantum codes.

2 Stabilizer Codes

2.1 Nondegenerate Stabilizer Codes

In the case of nondegenerate stabilizer codes just all bounds for classical codes are valid. For example we can use the best known asymptotic bound for nonbinary codes from [11]. (See figure 1, A). Note that as $\frac{\log_2 K}{n}$ tends to zero $\frac{d}{n} \approx 0.308$

2.2 General Case

Let Q be a quantum stabilizer $[[n, k]]$ code (degenerate or nondegenerate). In [7] it was shown that if the minimum distance of Q is d then there exist a classical binary code $[n-1, k, d]$. Using this observation and a classical linear programming bound, an upper bound for quantum stabilizer codes was

obtained [7] (see figure 1,C). Arguments used in [7] seem to be complicated. We show that a simpler approach yield to somewhat stronger bound.

Let C and $C^\perp, C \subseteq C^\perp$, be additive $[n, n-k]$ and $[n, n+k]$ codes over $\text{GF}(4)$ associated with Q . If an additive code has cardinality 2^t we will say that t is its dimension. So the dimensions of C and $C^\perp$ are $n-k$ and $n+k$ respectively. It is known that Q has minimum distance d iff all code words from the set $C^\perp \setminus C$ have weights greater than or equal to d (see [10] for details). We can consider codes C and $C^\perp$ as linear binary codes $\mathcal{C}$ and $\mathcal{C}^\perp$ of length $2n$. $\mathcal{C}^\perp$ is dual to $\mathcal{C}$ with respect to the symplectic inner product, i.e. if $(\mathbf{a}, \mathbf{b}) \in \mathcal{C}$ and $(\mathbf{a}', \mathbf{b}') \in \mathcal{C}^\perp$ then $(\mathbf{a}, \mathbf{b}) \bullet (\mathbf{a}', \mathbf{b}') = \langle \mathbf{a}, \mathbf{b}' \rangle \oplus \langle \mathbf{a}', \mathbf{b} \rangle = 0$. In [7] it was shown that a generator matrix G of $\mathcal{C}$ can be written in the form

$$G = \begin{bmatrix} I_s & A_1 & A_2 & B_1 & B_2 & 0 \\ 0 & 0 & 0 & D_1 & D_2 & I_r \end{bmatrix},$$

where I_s and B_1 are $s \times s$ matrices, A_1 and B_1 are $s \times k$ matrices, A_2 and B_2 are $s \times r$ matrices, D_1 is $r \times s$ matrix, and D_2 is $r \times k$ matrix. It is not difficult to check that all rows of the matrix $G^\perp$

$$G^\perp = \begin{bmatrix} I_s & A_1 & A_2 & B_1 & B_2 & 0 \\ 0 & 0 & 0 & D_1 & D_2 & I_r \\ 0 & 0 & 0 & A_1^\top & I_k & 0 \\ 0 & I_k & D_2^\top & B_2^\top & o & 0 \end{bmatrix},$$

are linearly independent and orthogonal (with respect to the symplectic inner product) to rows of the matrix G . So the matrix $G^\perp$ is a generator matrix for the code $\mathcal{C}^\perp$.

It is obvious that if the minimum distance of codewords from $C^\perp \setminus C$ is d then the minimum distance of the binary code, say S , with generator matrix

$$\begin{bmatrix} 0 & 0 & 0 & A_1^\top & I_k & 0 \\ 0 & I_k & D_2^\top & B_2^\top & o & 0 \end{bmatrix} \quad (6)$$

is not less than d . The code S is an $[n+k, 2k]$ binary code. Applying now any classical bound for this code we get a bound for code Q . For example we can use the second McEliece-Rodemich-Rumsey-Welch bound [8]. The resulting quantum bound is better than the bound from [7] (see figure 1, D). In fact in [7] only a code with the generator matrix

$$\begin{bmatrix} 0 & 0 & 0 & A_1^\top & I_k & 0 \end{bmatrix}$$

was taken into account.

We can get a stronger bound if we consider codes C and $C^\perp$ over $GF(4)$. A generator matrix of C can be written in the form [10]

$$G = \begin{bmatrix} I_{k_0} & \omega A_1 & B_1 \\ \omega I_{k_0} & \omega A_2 & B_2 \\ & I_{k_1} & A_3 \end{bmatrix}, \quad (7)$$

where B_j is a binary matrix, A_j is an arbitrary matrix, and ω is an element of $GF(4)$ of order 3. We will say that G defines a code of type $4^{k_0}2^{k_1}$. If in some l coordinates an additive code, say S , of length n contains only 0-s and α -s, $\alpha \in GF(4), \alpha \neq 0$, then we will say that S is a mixed code of lengths l and $n-l$. Note that if C is a code of type $4^{k_0}2^{k_1}$ of length n then corresponding quantum code Q is an $[[n, k]] = [[n, 2n-2k_0-k_1]]$ code.

Proposition 1 *Let d be a minimum distance of Q . Then d is not greater than the minimum distance of the best additive mixed code of lengths k_1 and $n-k_0-k_1$ and dimension $2n-4k_0-2k_1$. In particular*

- i) *if $k_1 = 0$ then d is not greater than the minimum distance of the best $[n-k_0, 2n-4k_0] = [(n+k)/2, 2k]$ additive code.*
- ii) *if $k_1 < \frac{2n-4k_0}{3} = 2k$ then d is not greater than the minimum distance of the best $[n-k_0-k_1, 2n-4k_0-3k_1] = [\frac{n+k-k_1}{2}, 2k-k_1]$ additive code.*

Proof

Let G' be a generator matrix of a complementary code of C to $C^\perp$, i.e. $\begin{bmatrix} G \\ G' \end{bmatrix}$ is a generator matrix of the code $C^\perp$. Due to the structure (7) of matrix G we can make elements of G' on the first k_0 positions equal 0 and elements on the next k_1 positions equal to 0 or ω . So G' will have the form $[0 \ D_1 \ D_2]$ where D_1 is an $2n-4k_0-2k_1 \times k_1$ matrix consisting from 0-s and ω -s and D_2 is an arbitrary $2n-4k_0-2k_1 \times n-k_0-k_1$ matrix. Since the minimum distance of a complementary code has to be not less than minimum distance of the quantum code Q we get the assertion.

i) follows from the previous.

ii) In the case $k_1 < \frac{2n-4k_0}{3}$ the matrix $[0 \ D_1 \ D_2]$ can be transformed to the form

$$\begin{bmatrix} 0 & A_1 & B_1 \\ 0 & 0 & B_2 \end{bmatrix},$$

where A_1 is an $k_1 \times k_1$ matrix consisting from 0-s and ω -s, B_1 and B_2 are arbitrary $k_1 \times n-k_0-k_1$ and $2n-4k_0-3k_1 \times n-k_0-k_1$ matrices. Since the

subcode with the generator matrix $[0 \ 0 \ B_2]$ has length $n - k_0 - k_1$, dimension $2n - 4k_0 - 3k_1$ and its minimum distance has to be not less than minimum distance of Q , the assertion follows. $\square$

Let S be a mixed code of lengths l and $n - l$ and dimension k .

Using standard arguments we can get the Hamming bound for the code S .

Lemma 1

$$\sum_{i=0}^e \sum_{j=0}^i \binom{l}{j} 3^{j-i} \binom{n-l}{i-j} \leq 2^{2n-l-k},$$

where $e = \left\lceil \frac{d-1}{2} \right\rceil$.

Combining Lemma 1 and Proposition 1 we get a bound of Hamming type for degenerate codes.

Proposition 2

$$\sum_{i=0}^e \sum_{j=0}^i \binom{k_1}{j} 3^{j-i} \binom{n-k_0-k_1}{i-j} \leq 2^{2k_0+3k_1} = 4^{\frac{n-k}{2}+k_1},$$

where $e = \left\lceil \frac{d-1}{2} \right\rceil$.

Consider the case $k_1 = 0$. Then we have

$$\sum_{i=0}^e 3^i \binom{n-k_0}{i} \leq 4^{\frac{n-k}{2}}.$$

Asymptotically this becomes

$$4^{\frac{n+k}{2}} H_4\left(\frac{d}{n+k}\right) \leq 4^{\frac{n-k}{2}}$$

$$\lambda = \frac{k}{n} \leq \frac{1 - H_4(\mu)}{1 + H_4(\mu)}; \quad \delta = \frac{d}{n} \geq \left(1 + \frac{1 - H_4(t)}{1 + H_4(t)}\right) t.$$

In the case $k_1 = 0$ we can also apply asymptotic bounds for complementary codes $[(n+k)/2, 2k]$ from Proposition 1. For example we can use an asymptotic bound from [11] (see fig. 1,E). Note that as $\frac{\log_2 K}{n}$ tends to zero $\frac{d}{n} = 0.375$.

Let $k_1 > 0$. Then according to Proposition 1 we can estimate a minimum distance of Q for $n - k_1 < k < k_1/2$ as minimum distance of an additive $[\frac{n+k-k_1}{2}, 2k - k_1]$ code. For example using the asymptotic bound from [11] and fixing $k_1 = 0, k_1 = 0.2, k_1 = 0.5$ we get bounds A,B,C, figure 2.

If $k > k_1/2$ we have to apply bounds for a mixed code. Generalizations of asymptotic bounds for mixed codes are not known to the author. Note that we can still use asymptotic bounds based on representation of complimentary code in the binary form (6) or the bound from [7] (bounds C and D on fig.1) for an arbitrary quantum stabilizer code.

References

- [1] P.W. Shor and R. Laflamme, "Quantum analog of the MacWilliams identities in classical coding theory," LANL e-print quant-ph/9610040.
- [2] E.M. Rains, "Quantum weight enumerators," LANL e-print quant-ph/9612015.
- [3] V.I. Levenshtein, "Krawtchouk Polynomials and Universal Bounds for Codes and Designs in Hamming Spaces," *IEEE Trans. on Inf. Theory*, **41**, no. 5, 1995, pp. 1303–1321.
- [4] P. Delsarte, "Bounds for unrestricted codes, by linear programming," *Philips Res. Reports*, **27**, 1972, pp. 272–289.
- [5] E. Knill and R. Laflamme, "A theory of quantum error correcting codes," LANL e-print quant-ph/9604034.
- [6] E.M. Rains, "Nonbinary quantum codes," LANL e-print quant-ph/97?.
- [7] R. Cleve, "Quantum stabilizer codes and classical linear codes," LANL e-print quant-ph/9612048.
- [8] R.J. McEliece, E.R. Rodemich, H. Rumsey Jr., and L.R. Welch, "New upper bounds on the rate of a code via the Delsarte-MacWilliams inequalities," *IEEE Trans. on Inf. Theory*, vol. IT-23, 1977, pp. 157–166.
- [9] E.M. Rains, "Quantum shadow enumerators," LANL e-print quant-ph/961101.

- [10] A.R. Calderbank, E.M. Rains, P.W. Shor and N.J.A. Sloane, “Quantum errors correction via codes over $\text{GF}(4)$,” LANL e-print quant-ph/9608006.
- [11] M. J. Aaltonen, “A new bound on nonbinary block codes,” *Discr. Math.*, vol. 83, pp. 139–160, 1990.




Figure 1: *A* is a bound for nondegenerate stabilizer codes; *B* is a bound for degenerate codes; *C* is a bound for degenerate stabilizer codes from [7]; *D* is bound based on a complementary code written in the binary form (6); *E* is a bound for degenerate stabilizer codes based on the classical bound from [11].




Figure 2: *A is a bound for $k_1 = 0$; B is a bound for $k_1 = 0.2$; C is a bound for $k_1 = 0.5$.*